arXiv:1506.01654v1 [math.AC] 4 Jun 2015

An inversion algorithm for polynomial maps

ELŻBIETA ADAMUS
Faculty of Applied Mathematics,
AGH University of Science and Technology
al. Mickiewicza 30, 30-059 Kraków, Poland
e-mail: esowa@agh.edu.pl

PAWEŁ BOGDAN
Faculty of Mathematics and Computer Science,
Jagiellonian University
ul. Łojasiewicza 6, 30-348 Kraków, Poland
e-mail: pawel.bogdan@uj.edu.pl

TERESA CRESPO
Departament d'Àlgebra i Geometria,
Universitat de Barcelona
Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain
e-mail: teresa.crespo@ub.edu

ZBIGNIEW HAJTO
Faculty of Mathematics and Computer Science,
Jagiellonian University
ul. Łojasiewicza 6, 30-348 Kraków, Poland
e-mail: zbigniew.hajto@uj.edu.pl

Abstract

We present an algorithmic equivalent statement to the Jacobian conjecture. Given a polynomial map F on an affine space of dimension n, our algorithm constructs n sequences of polynomials such that F is invertible if and only if the zero polynomial appears in all n sequences. This algorithm provides then a classification of polynomial automorphisms of affine spaces.

1 Introduction

The Jacobian Conjecture originated in the question raised by Keller in [8] on the invertibility of polynomial maps with Jacobian determinant equal to 1. The question is still open in spite of the efforts of many mathematicians. We recall in the sequel the precise statement of the Jacobian Conjecture, some reduction theorems and other results we shall use. We refer to [6] for a detailed account of the research on the Jacobian Conjecture and related topics.

Let K be a field and $K[X] = K[X_1, \dots, X_n]$ the polynomial ring in the variables $X_1, \dots, X_n$ over K. A *polynomial map* is a map $F = (F_1, \dots, F_n) : K^n \to K^n$ of the form

$$(X_1, \dots, X_n) \mapsto (F_1(X_1, \dots, X_n), \dots, F_n(X_1, \dots, X_n)),$$

where $F_i \in K[X], 1 \leq i \leq n$. The polynomial map F is *invertible* if there exists a polynomial map $G = (G_1, \dots, G_n) : K^n \to K^n$ such that $X_i = G_i(F_1, \dots, F_n), 1 \leq i \leq n$. We shall call F a *Keller map* if the Jacobian matrix

$$J = \left(\frac{\partial F_i}{\partial X_j} \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}}$$

has determinant equal to 1. Clearly an invertible polynomial map F has a Jacobian matrix J with non zero determinant and may be transformed into a Keller map by composition with the linear automorphism with matrix $J(0)^{-1}$.

Jacobian Conjecture. *Let K be a field of characteristic zero. A Keller map $F : K^n \to K^n$ is invertible.*

For $F = (F_1, \dots, F_n) \in K[X]^n$, we define the *degree* of F as $\deg F = \max\{\deg F_i : 1 \leq i \leq n\}$. It is known that if F is a polynomial automorphism, then $\deg F^{-1} \leq (\deg F)^{n-1}$ (see [1] or [9]).

The Jacobian conjecture for quadratic maps was proved by Wang in [10]. We state now the reduction of the Jacobian conjecture to the case of maps of third degree (see [1], [11], [4] and [5]).

Proposition 1. *a) (Bass-Connell-Wright-Yagzhev) Given a Keller map $F : \mathbb{C}^n \to \mathbb{C}^n$, there exists a Keller map $\widetilde{F} : \mathbb{C}^N \to \mathbb{C}^N$, $N \geq n$ of the form*

$\widetilde{F} = Id + H$, where $H(X)$ is a homogeneous cubic map and having the following property: if $\widetilde{F}$ is invertible, then F is invertible too.

b) (Drużkowski) The cubic part H may be chosen of the form

$$\left((\sum_{j=1}^{N} a_{1j} X_j)^3, \ldots, (\sum_{j=1}^{N} a_{Nj} X_j)^3 \right)$$

and with the matrix $A = (a_{ij})_{\substack{1 \le i \le N \\ 1 \le j \le N}}$ satisfying $A^2 = 0$.

Polynomial maps in the Drużkowski form are easier to handle than general cubic homogeneous polynomial maps. However we note the following result.

Proposition 2 ([7] Proposition 2.9)**.** *Let $r \in \mathbb{N}$. If the Jacobian Conjecture holds for all cubic homogeneous polynomial maps in r variables, then for all $n \in \mathbb{N}$ the Jacobian Conjecture holds for all polynomial maps of the form*

$$F = X + (AX)^3$$

with $A \in \mathrm{M}_n(\mathbb{C})$ and $\operatorname{rank} A \le r$.

Given a polynomial map $F : \mathbb{C}^n \to \mathbb{C}^n$, we shall call a polynomial $P \in \mathbb{C}[X_1, \ldots, X_n]$ *invariant under F* if $P(F_1, \ldots, F_n) = P(X_1 \ldots, X_n)$. A polynomial map $F : \mathbb{C}^n \to \mathbb{C}^n$ of the form $F = Id + H$ is called a *quasi-translation* if $F^{-1} = Id - H$.

In this paper we present a recursive algorithm to invert polynomial maps. Given a polynomial map $F : \mathbb{C}^n \to \mathbb{C}^n$ of the form $F = Id + H$, where $H(X)$ is a homogeneous cubic map, our algorithm constructs, for $1 \le i \le n$, a sequence P^i_k of polynomials in $\mathbb{C}[X]$ with $P^i_0 = X_i$ such that F is invertible if and only $P^i_{m_i} = 0$, for some integer $m_i > 0$, $1 \le i \le n$ and, when this is the case, we may compute F^{-1} as an alternating sum of the polynomials P^i_k, $0 \le k < m_i$. In the last section, we apply the algorithm to several examples of polynomial maps.

2 The algorithm

Let us consider a polynomial map $F : \mathbb{C}^n \to \mathbb{C}^n$. Given a polynomial $P(X_1, \ldots, X_n) \in \mathbb{C}[X] = \mathbb{C}[X_1, \ldots, X_n]$, we define the following sequence of polynomials in $\mathbb{C}[X]$,

$$\begin{aligned} P_0(X_1,\dots,X_n) &= P(X_1,\dots,X_n), \\ P_1(X_1,\dots,X_n) &= P_0(F_1,\dots,F_n) - P_0(X_1,\dots,X_n), \end{aligned}$$

and, assuming P_{k-1} is defined,

$$P_k(X_1,\dots,X_n) = P_{k-1}(F_1,\dots,F_n) - P_{k-1}(X_1,\dots,X_n).$$

The following lemma is easy to prove.

Lemma 3. *For a positive integer m, we have*

$$P(X_1,\dots,X_n) = \sum_{l=0}^{m-1} (-1)^l P_l(F_1,\dots,F_n) + (-1)^m P_m(X_1,\dots,X_n).$$

In particular, if we assume that for some integer m, $P_m(X_1,\dots,X_n) = 0$, then

$$P(X_1,\dots,X_n) = \sum_{l=0}^{m-1} (-1)^l P_l(F_1,\dots,F_n).$$

Theorem 4. *Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a polynomial map of the form*

$$\begin{cases} F_1(X_1,\dots,X_n) &= X_1 + H_1(X_1,\dots,X_n) \\ F_2(X_1,\dots,X_n) &= X_2 + H_2(X_1,\dots,X_n) \\ &\vdots \\ F_n(X_1,\dots,X_n) &= X_n + H_n(X_1,\dots,X_n), \end{cases}$$

where $H_i(X_1,\dots,X_n)$ is a homogeneous polynomial in $X_1,\dots,X_n$ of degree 3, $1 \le i \le n$. Let us assume that F is invertible. Then for the polynomial sequence (P_k^i) constructed with $P = X_i$, there exists an integer m_i such that $P_{m_i}^i = 0$ and the inverse map G of F is given by

$$G_i(Y_1,Y_2,\dots,Y_n) = \sum_{l=0}^{m_i-1} (-1)^l P_l^i(Y_1,Y_2,\dots,Y_n),\ 1 \le i \le n.$$

Proof. Taking into account lemma 3, it is enough to prove that there exists an integer m_i such that $P_{m_i}^i = 0$, for $i = 1,\dots,n$. If F is invertible, then $G = F^{-1}$ is of the form

$$\begin{cases} G_1(Y_1, Y_2, \ldots, Y_n) & = & Y_1 + J_1 \\ G_2(Y_1, Y_2, \ldots, Y_n) & = & Y_2 + J_2 \\ & \vdots & \\ G_n(Y_1, Y_2, \ldots, Y_n) & = & Y_n + J_n, \end{cases}$$

where J_i is a polynomial in $Y_1, \ldots, Y_n$, $1 \leq i \leq n$. Let $N_i = \deg J_i$. Let us consider, for a fixed i, the polynomial sequence

$$\begin{array}{rcl} P_0^i(X_1, \ldots, X_n) & = & X_i, \\ P_1^i(X_1, \ldots, X_n) & = & F_i(X_1, \ldots, X_n) - X_i = H_i(X_1, \ldots, X_n), \\ P_2^i(X_1, \ldots, X_n) & = & H_i(F_1, \ldots, F_n) - H_i(X_1, \ldots, X_n), \\ & \vdots & \end{array}$$

We write the Taylor series for the polynomial $H_i(F_1, \ldots, F_n) = H_i(X_1 + H_1, \ldots, X_n + H_n)$ and obtain

$$\begin{array}{rcl} P_2(X_1, \ldots, X_n) & = & H_i(F_1, \ldots, F_n) - H_i(X_1, \ldots, X_n) \\ & = & Q_{21} + Q_{22} + Q_{23} \end{array}$$

where

$$\begin{array}{rcl} Q_{21} & = & \sum_{j=1}^n H_j \dfrac{\partial H_i}{\partial X_j} \\ Q_{22} & = & \dfrac{1}{2} \sum_{1 \leq j,k \leq n} H_j H_k \dfrac{\partial^2 H_i}{\partial X_j \partial X_k} \\ Q_{23} & = & \dfrac{1}{6} \sum_{1 \leq j,k,l \leq n} H_j H_k H_l \dfrac{\partial^3 H_i}{\partial X_j \partial X_k \partial X_l} \end{array}$$

and the polynomials Q_{21}, Q_{22}, Q_{23} are either zero or homogeneous polynomials of degrees $5, 7, 9$, respectively. Let us prove by induction that

$$P_n^i = \sum_{j=1}^{(3^n - 2n + 1)/2} Q_{nj},$$

where Q_{nj} is either zero or a homogeneous polynomial of degree $2n - 1 + 2j$. We have already seen it for $n = 2$. Let us assume $P_{n-1}^i = \sum_{j=1}^{(3^{n-1} - 2n + 3)/2} Q_{n-1,j}$ with $Q_{n-1,j}$ either zero or a homogeneous polynomial of degree $2n - 3 + 2j$. We want to prove the property for P_n^i. We have

$$P_n^i(X_1,\dots,X_n) = P_{n-1}^i(F_1,\dots,F_n) - P_{n-1}^i(X_1,\dots,X_n)$$
$$= \textstyle\sum_{j=1}^{(3^{n-1}-2n-1)/2} Q_{n-1,j}(F_1,\dots,F_n) - Q_{n-1,j}(X_1,\dots,X_n).$$

If $Q(X_1,\dots,X_n)$ is a homogeneous polynomial of degree d,

$$Q(F_1,\dots,F_n) - Q(X_1,\dots,X_n) = \sum_{j=1}^{n} H_j \frac{\partial Q}{\partial X_j} + \dots$$

is a sum of polynomials which are either zero or homogeneous polynomials of degrees $3+d-1, 6+d-2, \dots, 3d$. Hence $Q_{n-1,j}(F_1,\dots,F_n) - Q_{n-1,j}(X_1,\dots,X_n)$ is a sum of polynomials which are either zero or homogeneous polynomials of degrees $2n+2j-1, 2n+2j+1, \dots, 6n-9+6j$. Therefore $P_n^i = \sum_{j=1}^{(3^n-2n+1)/2} Q_{nj}$ where Q_{nj} is either zero or a homogeneous polynomial of degree $2n-1+2j$.

Applying lemma 3, we obtain

$$X_i = \sum_{l=0}^{m-1} (-1)^l P_l(F_1,\dots,F_n) + (-1)^m P_m(X_1,\dots,X_n).$$

Now, we have that $X_i = G_i(F_1,\dots,F_n)$ and G_i is a polynomial of degree N_i. Then the summands appearing in the development of the expression $G_i(F_1(X_1,\dots,X_n),\dots,F_n(X_1,\dots,X_n))$ have total degree $3N_i$ in $X_1,\dots,X_n$. If $m > (3N_i-1)/2$, then P_m is a sum of polynomials which are either zero or homogeneous of degrees bigger than $3N_i$. Hence $P_m = 0$. $\square$

We shall give now a bound for the degrees of the polynomials P_k^i in Theorem 4.

Proposition 5. *Let* F, G *and* (P_k^i) *be as in Theorem 4. Let* $N_i = \deg G_i$. *Then* $\deg P_k^i \le 3N_i$. *In particular, if* $P_{(3N_i-1)/2} \neq 0$, *then it is homogeneous of degree* $3N_i$.

Proof. We have seen in the proof of Theorem 4 that $\deg P_k^i \le 3N_i$, for $k \le [log_3 N_i] + 1$. Let $m = [log_3 N_i] + 2$. Applying Lemma 3, we obtain

$$X_i = \sum_{l=0}^{m-1} (-1)^l P_l^i(F_1, F_2, \dots, F_n) + (-1)^m P_m^i(X_1,\dots,X_n).$$

Reasoning as in the proof of Theorem 4, we obtain that the homogeneous summands of P^i_m of degrees $> 3N_i$ must be zero, hence $\deg P^i_m \leq 3N_i$. By induction we obtain $\deg P^i_k \leq 3N_i$ for all k. By Theorem 4, the homogeneous summands of $P_{(3N_i-1)/2}$ have degrees $\geq 3N_i$, so we obtain the last sentence of the Proposition. □

Corollary 6. *Let F be a polynomial map as in Theorem 4. If F is invertible, then there exists $P \in \mathbb{C}[X_1, \ldots, X_n]$ invariant under F.*

Proof. Indeed, if P^i_m is the first zero polynomial in the sequence (P^i_k), then P^i_{m-1} is invariant under F. □

Remark 7. We have presented our algorithm for cubic homogeneous complex polynomial maps but one can see easily that it is valid for cubic homogeneous polynomial maps defined over any field K of characteristic 0.

Due to Proposition 1, we have presented the algorithm in the cubic homogeneous case however its validity is wider.

Proposition 8. *Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a polynomial map of the form*

$$\begin{cases} F_1(X_1, \ldots, X_n) &= X_1 + H_1(X_1, \ldots, X_n) \\ F_2(X_1, \ldots, X_n) &= X_2 + H_2(X_1, \ldots, X_n) \\ & \vdots \\ F_n(X_1, \ldots, X_n) &= X_n + H_n(X_1, \ldots, X_n), \end{cases}$$

where $H_i(X_1, \ldots, X_n)$ is a polynomial in $X_1, \ldots, X_n$ of lower degree $d_i \geq 2$, $1 \leq i \leq n$. Let us assume that F is invertible. Then for the polynomial sequence (P^i_k) constructed with $P = X_i$, there exists an integer m_i such that $P^i_{m_i} = 0$ and the inverse map G of F is given by

$$G_i(Y_1, Y_2, \ldots, Y_n) = \sum_{l=0}^{m_i-1} (-1)^l P^i_l(Y_1, Y_2, \ldots, Y_n),\ 1 \leq i \leq n.$$

Proof. Following the proof of Theorem 4, one can see that the lower degrees of the polynomials (P^i_k) are increasing by (at least) $d - 1 > 0$, where $d = \min\{d_i\}$, hence we obtain $P^i_m = 0$, for $m > (DN_i - d_i)/(d-1) + 1$, where $D = \deg F, N_i = \deg G_i$. □

As a consequence of Proposition 8, we obtain a direct proof of the characterization of quasi-translations obtained by de Bondt in [2], Proposition 1.1. See also [3] Proposition 1.3.

Proposition 9. *Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a polynomial map of the form $F = Id + H$, where $H_i(X_1, \dots, X_n)$ is a polynomial in $X_1, \dots, X_n$ of lower degree $d_i \geq 2$, $1 \leq i \leq n$. Then the following conditions are equivalent.*

a) F is a quasi-translation.

b) $JH \cdot H = 0$, where JH denotes the jacobian matrix of H.

c) $P_2^i = 0$, $1 \leq i \leq n$, where (P_k^i) denotes the polynomial sequence constructed with $P = X_i$.

Proof. From Lemma 3 applied to $P(X) = X_i$, we have $X_i = F_i - H_i(F) + P_2^i(X)$, hence the equivalence between a) and c) is clear. Now

$$P_2^i(X) = H_i(F) - H_i(X) = \sum_{k \geq 1} \sum_{1 \leq j_1, \dots, j_k \leq n} \frac{\partial^k H_i}{\partial X_{j_1} \dots \partial X_{j_k}} H_{j_1} \dots H_{j_k}.$$

Let us assume b). We have then $\sum_{j=1}^n \dfrac{\partial H_i}{\partial X_j} H_j = 0$, $1 \leq i \leq n$. We shall prove by induction

$$\sum_{1 \leq j_1, \dots, j_k \leq n} \frac{\partial^k H_i}{\partial X_{j_1} \dots \partial X_{j_k}} H_{j_1} \dots H_{j_k} = 0 \tag{1}$$

for all $k \geq 0$. For $k = 1$, it is true by hypothesis. Let us assume that it is true for k. By applying $\partial / \partial X_j$ to (1), we obtain

$$\sum_{1 \leq j_1, \dots, j_k \leq n} \left(\frac{\partial^{k+1} H_i}{\partial X_{j_1} \dots \partial X_{j_k} \partial X_j} H_{j_1} \dots H_{j_k} + \sum_{l=1}^{k} \frac{\partial^k H_i}{\partial X_{j_1} \dots \partial X_{j_k}} H_{j_1} \dots \frac{\partial H_{j_l}}{\partial X_j} \dots H_{j_k} \right) = 0.$$

Multiplying by H_j, summing from $j = 1$ to n and taking into account the case $k = 1$ with $i = j_1, \dots, j_k$, we obtain (1) for $k + 1$.

Let us assume c). If H is homogeneous, the summands of P_2^i are homogeneous of different degrees, hence all of them vanish and since the first summand $\sum_{j=1}^n \dfrac{\partial H_i}{\partial X_j} H_j$ is the i-th row of $JH \cdot H$, we obtain b). If H is not homogeneous, we obtain successively the vanishing of the homogeneous summands of $\sum_{j=1}^n \dfrac{\partial H_i}{\partial X_j} H_j$ in increasing degree order. □

Remark 10. In the set of polynomial maps as in Proposition 9, our algorithm provides a filtration $\{\mathcal{P}_n\}_{n\geq 1}$ such that $\mathcal{P}_n$ is the subset of polynomial maps with $P_n^i = 0, 1 \leq i \leq n$. Clearly $\mathcal{P}_1 = \{Id\}$ and, applying Proposition 9, $\mathcal{P}_2$ is the set of quasi-translations.

3 Examples

3.1

We consider the following Keller map in dimension 2 to illustrate that the validity of our algorithm is not restricted to homogeneous cubic maps.

$$\left\{ \begin{array}{lcl} F_1 & = & X_1 + (X_2 + X_1^3)^2 \\ F_2 & = & X_2 + X_1^3 \end{array} \right.$$

Let us write $H_1 := (X_2 + X_1^3)^2, H_2 := X_1^3$. Taking into account the bound for the degree of the inverse, we may disregard the terms of degree > 6 in the polynomial sequences (P_k^i). We obtain

$$\begin{array}{lcl} P_0^1 & = & X_1 \\ P_1^1 & = & H_1 = (X_2 + X_1^3)^2 \\ P_2^1 & = & 3X_1^6 + 6X_1X_2^5 + 6X_1^2X_2^3 + 2X_1^3X_2 \\ P_3^1 & = & 2X_1^6 + 18X_1X_2^5 + 6X_1^2X_2^3 \\ P_4^1 & = & 12X_1X_2^5 \\ P_5^1 & = & 0 \end{array}$$

which gives $X_1 = P_0^1(F_1, F_2) - P_1^1(F_1, F_2) + P_2^1(F_1, F_2) - P_3^1(F_1, F_2) + P_4^1(F_1, F_2) = Y_1 - Y_2^2$ and

$$\begin{array}{lcl} P_0^2 & = & X_2 \\ P_1^2 & = & H_2 = X_1^3 \\ P_2^2 & = & 6X_1^5X_2 + X_2^6 + 3X_1X_2^4 + 3X_1^2X_2^2 \\ P_3^2 & = & 6X_1^5X_2 + 6X_2^6 + 6X_1X_2^4 \\ P_4^2 & = & 6X_2^6 \\ P_5^2 & = & 0 \end{array}$$

which gives $X_2 = P_0^2(F_1, F_2) - P_1^2(F_1, F_2) + P_2^2(F_1, F_2) - P_3^2(F_1, F_2) + P_4^2(F_1, F_2) = Y_2 - Y_1^3 + Y_2^6 - 3Y_1Y_2^4 + 3Y_1^2Y_2^2 = Y_2 - (Y_1 - Y_2^2)^3$.

3.2

We consider the following Keller map F in dimension 5.

$$\left\{\begin{array}{rcl}
F_1 & = & X_1 + a_1X_4^3 + a_2x_4^2X_5 + a_3X_4X_5^2 + a_4x_5^3 + \dfrac{a_2c_5X_2X_4^2}{c_2} + \dfrac{2a_3c_5X_2X_4X_5}{c_2} \\
 & & + \dfrac{3a_4c_5X_2X_5^2}{c_2} - \dfrac{a_2e_2X_3X_4^2}{c_2} - \dfrac{2a_3e_2X_3X_4X_5}{c_2} - \dfrac{3a_4e_2X_3X_5^2}{c_2} \\
 & & + \dfrac{a_3c_5^2X_2^2X_4}{c_2^2} + \dfrac{3a_4c_5^2X_2^2X_5}{c_2^2} - \dfrac{2a_3c_5e_2X_2X_3X_4}{c_2^2} - \dfrac{6a_4c_5e_2X_2X_3X_5}{c_2^2} \\
 & & + \dfrac{a_3e_2^2X_3^2X_4}{c_2^2} + \dfrac{3a_4e_2^2X_3^3X_5}{c_2^2} + \dfrac{a_4e_5^3X_2^3}{c_2^3} - \dfrac{3a_4c_5^2e_2X_2^2X_3}{c_2^3} \\
 & & + \dfrac{3a_4c_5e_2^2X_2X_3^2}{c_2^3} - \dfrac{a_4e_2^3X_3^3}{c_2^3} \\
F_2 & = & X_2 + b_1X_4^3 \\
F_3 & = & X_3 + c_5X_2X_4^2 + c_1X_4^3 + c_2X_4^4X_5 - e_2X_3X_4^2 \\
F_4 & = & X_4 \\
F_5 & = & X_5 + e_2X_4^2X_5 + \dfrac{c_5e_2X_2X_4^2}{c_2} - \dfrac{e_2^2X_3X_4^2}{c_2} - \dfrac{(b_1c_5 - c_1e_2)X_4^3}{c_2}
\end{array}\right.$$

with parameters $a_1, a_2, a_3, a_4, b_1, c_1, c_2, c_5, e_2$. By applying the algorithm we obtain $P_2^i = 0$, for $i = 1, \ldots, 5$, hence F is a quasi-translation.

3.3

We consider the following Keller map F in dimension 6

$$\left\{\begin{array}{rcl}
F_1 & = & X_1 + a_5e_1(X_1 + X_2)^3/a_4 + a_4X_2X_4X_6 + a_5X_4X_5X_6 \\
F_2 & = & X_2 - a_5e_1(X_1 + X_2)^3/a_4 \\
F_3 & = & X_3 + c_1X_1^3 + c_2(X_1 + X_5)^3 + c_3(X_1 + X_2)^3 + c_4(X_1 + X_4)^3 + c_5X_6^3 \\
F_4 & = & X_4 + d_4X_2X_6^2 + a_5d_4X_5X_6^2/a_4 \\
F_5 & = & X_5 + e_1(X_1 + X_2)^3 \\
F_6 & = & X_6
\end{array}\right.$$

with parameters $a_4, a_5, c_1, c_2, c_3, c_4, c_5, d_4, e_1$. Denoting $G = F^{-1}$ and taking variables $(Y_1, \ldots, Y_6)$ for G, we obtain $P_8^1 = 0$, $P_9^2 = 0$ and

$$
\begin{aligned}
&G_1 = -(20a_5^4e_1Y_6^9d_4^3Y_2^3Y_5^3a_4^3 - 6a_5^4e_1Y_6^4Y_4Y_5^3d_4Y_2a_4^2 + a_5e_1a_4^6Y_6^9Y_2^6d_4^3 + 3Y_6^3a_5e_1Y_2^4d_4a_4^4 \\
&+3a_5e_1a_4^5Y_6^6Y_2^5d_4^2 - a_5e_1a_4^6Y_2^3Y_4^3Y_6^3 + a_4^5Y_2Y_4Y_6 - Y_6^3a_4^5d_4Y_2^2 + a_5^7e_1Y_6^9d_4^3Y_5^6 \\
&+a_5e_1a_4^3Y_1^3 + a_5e_1a_4^3Y_2^3 - Y_1a_4^4 - 3a_5^2e_1a_4^5Y_2^2Y_4^3Y_6^3Y_5 - 6a_5e_1Y_1Y_2^2Y_4Y_6a_4^4 \\
&+3a_5^5e_1Y_6^6Y_1d_4^2Y_5^4a_4 + 12a_5^4e_1Y_6^6Y_2^2d_4^2Y_5^3a_4^2 - 6a_5^4e_1Y_6^4Y_1Y_4d_4Y_5^3a_4^2 + 3a_5e_1a_4^3Y_1Y_2^2 \\
&-6a_5e_1a_4^5Y_6^4Y_1Y_2^3Y_4d_4 + 3Y_6^3a_5e_1Y_1^2Y_2^2d_4a_4^4 + 3a_5e_1a_4^5Y_1Y_2^2Y_4^2Y_6^2 - 30a_5^3e_1Y_6^7Y_4d_4^2Y_2^3Y_5^2a_4^4 \\
&+6Y_6^3a_5e_1Y_1Y_2^3d_4a_4^4 + 15a_5^3e_1Y_6^9d_4^3Y_2^4Y_5^2a_4^4 + 6a_5^2e_1a_4^5Y_6^9Y_2^5d_4^3Y_5 - 18a_5^2e_1Y_6^4Y_2^3Y_4d_4Y_5a_4^4 \\
&+12a_5^2e_1a_4^5Y_6^5Y_2^3Y_4^2d_4Y_5 - 15a_5^2e_1a_4^5Y_6^7Y_2^4Y_4d_4^2Y_5 + 12a_5^2e_1Y_6^6Y_1Y_2^3d_4^2Y_5a_4^4 - 18a_5^2e_1Y_6^4Y_1Y_2^2Y_4d_4Y_5a_4^4 \\
&-3a_5e_1a_4^6Y_6^7Y_2^5Y_4d_4^2 + 3a_5e_1a_4^3Y_1^2Y_2 + 18a_5^3e_1Y_6^5Y_4^2d_4Y_2^2Y_5^2a_4^4 + 12a_5^2e_1Y_6^6Y_2^4d_4^2Y_5a_4^4 \\
&+3a_5e_1a_4^5Y_6^6Y_1Y_2^4d_4^2 + 3a_5e_1a_4^6Y_6^5Y_2^4Y_4^2d_4 - 6a_5e_1Y_6^4a_4^5Y_2^4Y_4d_4 + a_5Y_4Y_5Y_6a_4^4 \\
&+15a_5^5e_1Y_6^9d_4^3Y_2^2Y_5^4a_4^2 + 6a_5^6e_1Y_6^9Y_2d_4^3Y_5^5a_4 - 18a_5^3e_1Y_6^4Y_2^2Y_4d_4Y_5^2a_4^3 + 18a_5^3e_1Y_6^6Y_1d_4^2Y_2^2Y_5^2a_4^3 \\
&-3a_5^6e_1Y_6^7Y_4Y_5^5d_4^2a_4 - 18a_5^3e_1Y_6^4Y_1Y_4d_4Y_2Y_5^2a_4^3 + 6a_5^3e_1Y_6^3Y_1d_4Y_5^2Y_2a_4^2 - 30a_5^4e_1Y_6^7Y_4d_4^2Y_2^2Y_5^3a_4^3 \\
&+12a_5^4e_1Y_6^5Y_2Y_4^2d_4Y_5^3a_4^3 + 12a_5^4e_1Y_6^6Y_1d_4^2Y_2Y_5^3a_4^2 + 3a_5^5e_1Y_6^6d_4^2Y_5^4Y_2a_4 + 3a_5^3e_1Y_6^3Y_1^2d_4Y_5^2a_4^2 \\
&+18a_5^3e_1Y_6^6Y_2^3d_4^2Y_5^2a_4^3 + 6a_5^2e_1Y_1Y_2Y_4^2Y_6^2Y_5a_4^4 - 3a_5e_1Y_2^3Y_4Y_6a_4^4 - 2Y_6^3Y_2a_5d_4Y_5a_4^4 \\
&+3a_5^5e_1Y_6^5Y_4^2Y_5^4d_4a_4^2 - 3a_5^3e_1Y_2Y_4^3Y_6^3Y_5^2a_4^4 - 3a_5e_1Y_1^2Y_2Y_4Y_6a_4^4 + 3a_5e_1a_4^5Y_2^3Y_4^2Y_6^2 \\
&+6a_5^2e_1Y_2^2Y_4^2Y_6^2Y_5a_4^4 - 3a_5^2e_1Y_4Y_5Y_6Y_2^2a_4^3 - a_5^2Y_6^3d_4Y_5^2a_4^3 - 3a_5^2e_1Y_1^2Y_4Y_5Y_6a_4^3 \\
&+3a_5^3e_1Y_4^2Y_5^2Y_6^2Y_2a_4^3 + 6Y_6^3a_5^2e_1Y_1^2Y_2d_4Y_5a_4^3 - 15a_5^5e_1Y_6^7Y_2Y_4d_4^2Y_5^4a_4^2 + 12Y_6^3a_5^2e_1Y_1Y_2^2d_4Y_5a_4^3 \\
&+6Y_6^3a_5^2e_1Y_2^3d_4Y_5a_4^3 + 3a_5^3e_1Y_6^3d_4Y_5^2Y_2^2a_4^2 + 3a_5^3e_1Y_1Y_4^2Y_5^2Y_6^2a_4^3 - 6a_5^2e_1Y_1Y_4Y_5Y_6Y_2a_4^3 \\
&-a_5^4e_1Y_4^3Y_5^3Y_6^3a_4^3)/a_4^4;
\end{aligned}
$$

$$
\begin{aligned}
&G_2 = (20a_5^4e_1Y_6^9d_4^3Y_2^3Y_5^3a_4^3 - 6a_5^4e_1Y_6^4Y_4Y_5^3d_4Y_2a_4^2 + a_5e_1a_4^6Y_6^9Y_2^6d_4^3 + 3Y_6^3a_5e_1Y_2^4d_4a_4^4 \\
&+3a_5e_1a_4^5Y_6^6Y_2^5d_4^2 - a_5e_1a_4^6Y_2^3Y_4^3Y_6^3 + a_5^7e_1Y_6^9d_4^3Y_5^6 + a_5e_1a_4^3Y_1^3 \\
&+a_5e_1a_4^3Y_2^3 - 3a_5^2e_1a_4^5Y_2^2Y_4^3Y_6^3Y_5 - 6a_5e_1Y_1Y_2^2Y_4Y_6a_4^4 + 3a_5^5e_1Y_6^6Y_1d_4^2Y_5^4a_4 \\
&+12a_5^4e_1Y_6^6Y_2^2d_4^2Y_5^3a_4^2 - 6a_5^4e_1Y_6^4Y_1Y_4d_4Y_5^3a_4^2 + 3a_5e_1a_4^3Y_1Y_2^2 - 6a_5e_1a_4^5Y_6^4Y_1Y_2^3Y_4d_4 \\
&+3Y_6^3a_5e_1Y_1^2Y_2^2d_4a_4^4 + 3a_5e_1a_4^5Y_1Y_2^2Y_4^2Y_6^2 - 30a_5^3e_1Y_6^7Y_4d_4^2Y_2^3Y_5^2a_4^4 + 6Y_6^3a_5e_1Y_1Y_2^3d_4a_4^4 \\
&+15a_5^3e_1Y_6^9d_4^3Y_2^4Y_5^2a_4^4 + 6a_5^2e_1a_4^5Y_6^9Y_2^5d_4^3Y_5 - 18a_5^2e_1Y_6^4Y_2^3Y_4d_4Y_5a_4^4 + 12a_5^2e_1a_4^5Y_6^5Y_2^3Y_4^2d_4Y_5 \\
&-15a_5^2e_1a_4^5Y_6^7Y_2^4Y_4d_4^2Y_5 + 12a_5^2e_1Y_6^6Y_1Y_2^3d_4^2Y_5a_4^4 - 18a_5^2e_1Y_6^4Y_1Y_2^2Y_4d_4Y_5a_4^4 - 3a_5e_1a_4^6Y_6^7Y_2^5Y_4d_4^2 \\
&+3a_5e_1a_4^3Y_1^2Y_2 + 18a_5^3e_1Y_6^5Y_4^2d_4Y_2^2Y_5^2a_4^4 + 12a_5^2e_1Y_6^6Y_2^4d_4^2Y_5a_4^4 + 3a_5e_1a_4^5Y_6^6Y_1Y_2^4d_4^2 \\
&+3a_5e_1a_4^6Y_6^5Y_2^4Y_4^2d_4 - 6a_5e_1Y_6^4a_4^5Y_2^4Y_4d_4 + 15a_5^5e_1Y_6^9d_4^3Y_2^2Y_5^4a_4^2 + 6a_5^6e_1Y_6^9Y_2d_4^3Y_5^5a_4 \\
&-18a_5^3e_1Y_6^4Y_2^2Y_4d_4Y_5^2a_4^3 + 18a_5^3e_1Y_6^6Y_1d_4^2Y_2^2Y_5^2a_4^3 - 3a_5^6e_1Y_6^7Y_4Y_5^5d_4^2a_4 - 18a_5^3e_1Y_6^4Y_1Y_4d_4Y_2Y_5^2a_4^3 \\
&+6a_5^3e_1Y_6^3Y_1d_4Y_5^2Y_2a_4^2 - 30a_5^4e_1Y_6^7Y_4d_4^2Y_2^2Y_5^3a_4^3 + 12a_5^4e_1Y_6^5Y_2Y_4^2d_4Y_5^3a_4^3 + 12a_5^4e_1Y_6^6Y_1d_4^2Y_2Y_5^3a_4^2 \\
&+3a_5^5e_1Y_6^6d_4^2Y_5^4Y_2a_4 + 3a_5^3e_1Y_6^3Y_1^2d_4Y_5^2a_4^2 + 18a_5^3e_1Y_6^6Y_2^3d_4^2Y_5^2a_4^3 + 6a_5^2e_1Y_1Y_2Y_4^2Y_6^2Y_5a_4^4 \\
&-3a_5e_1Y_2^3Y_4Y_6a_4^4 + 3a_5^5e_1Y_6^5Y_4^2Y_5^4d_4a_4^2 - 3a_5^3e_1Y_2Y_4^3Y_6^3Y_5^2a_4^4 - 3a_5e_1Y_1^2Y_2Y_4Y_6a_4^4 \\
&+3a_5e_1a_4^5Y_2^3Y_4^2Y_6^2 + 6a_5^2e_1Y_2^2Y_4^2Y_6^2Y_5a_4^4 - 3a_5^2e_1Y_4Y_5Y_6Y_2^2a_4^3 - 3a_5^2e_1Y_1^2Y_4Y_5Y_6a_4^3 \\
&+3a_5^3e_1Y_4^2Y_5^2Y_6^2Y_2a_4^3 + 6Y_6^3a_5^2e_1Y_1^2Y_2d_4Y_5a_4^3 - 15a_5^5e_1Y_6^7Y_2Y_4d_4^2Y_5^4a_4^2 + 12Y_6^3a_5^2e_1Y_1Y_2^2d_4Y_5a_4^3 \\
&+6Y_6^3a_5^2e_1Y_2^3d_4Y_5a_4^3 + 3a_5^3e_1Y_6^3d_4Y_5^2Y_2^2a_4^2 + 3a_5^3e_1Y_1Y_4^2Y_5^2Y_6^2a_4^3 - 6a_5^2e_1Y_1Y_4Y_5Y_6Y_2a_4^3 \\
&-a_5^4e_1Y_4^3Y_5^3Y_6^3a_4^3 + Y_2a_4^4)/a_4^4.
\end{aligned}
$$

Now,

$$
\begin{aligned}
G_6 &= Y_6 \\
G_5 &= Y_5 - e_1(G_1 + G_2)^3 \\
G_4 &= Y_4 - d_4G_2G_6^2 - a_5d_4G_5G_6^2/a_4 \\
G_3 &= Y_3 - c_1G_1^3 - c_2(G_1 + G_5)^3 - c_3(G_1 + G_2)^3 - c_4(G_1 + G_4)^3 + c_5G_6^3.
\end{aligned}
$$

Acknowledgments. E. Adamus acknowledges support of the Polish Ministry of Science and Higher Education. T. Crespo and Z. Hajto acknowledge support of grant MTM2012-33830, Spanish Science Ministry.

References

[1] H. Bass, E. Connell and D. Wright, *The Jacobian Conjecture: Reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. 7 (1982), 287-330.

[2] M. de Bondt, *Quasi-translations and counterexamples to the homogeneous dependence problem*, Proc. Amer. Math. Soc. 134 (2006), 2849-2856.

[3] M. de Bondt, *Quasi-translations and singular Hessians*, arXiv:1501.05168 [math.AG].

[4] L. Drużkowski, *An effective approach to Keller's Jacobian conjecture*, Math. Ann. 264 (1983), 303-313.

[5] L. Drużkowski, *New reduction in the Jacobian conjecture*, Univ. Iagel. Acta Math. No. 39 (2001), 203-206.

[6] A. van der Essen, *Polynomial automorphisms and the Jacobian Conjecture*, Progress in Mathematics 190, Birkhäuser Verlag, 2000.

[7] A. van der Essen, *Seven lectures on polynomial automorphisms*, Automorphisms of affine spaces (Curaçao, 1994), Kluwer Acad. Publ., Dordrecht, 1995, 3-39.

[8] O. H. Keller, *Ganze Cremona-Transformationen*, Monatsh. Math. Phys. 47 (1939), 299-306.

[9] K. Rusek and T. Winiarski, *Polynomial automorphisms of* $\mathbb{C}^n$, Univ. Iagell. Acta Math. 24 (1984), 143-149.

[10] S. Wang, *A Jacobian criterion for separability*, J. Algebra 65 (1980), 453-494.

[11] A. V. Yagzhev, *On a problem of O.H. Keller*, Sibirsk. Mat. Zh. 21 (1980), 747-754.